
Appendix D. McCown's Longspur Status and Trends.

Status and trends are summarized and organized at three geographic scales that are utilized in analyses by the Breeding Bird Survey (BBS): Survey wide/country, Bird Conservation Region (BCR), and state/province. Official Species Status (Regulatory): Species at Risk Act (SARA) and State status: "T" = Threatened. State/Provincial Conservation Status represents State Wildlife Action Plan (SWAP) status and Provincial status: Tiers range 1-3 with Tier 1 the highest level of conservation priority. "SGCN" (Species of Greatest Conservation Need) are priority species without assigned tiers. Tier 1C and S4 are low designations of conservation priority for Arizona and Texas, respectively. For

the BBS trends: n = number of survey routes on which the species was encountered during the entire (1967–2015) interval. BBS trends are presented as yearly percentage change. Numbers in parentheses are the credible intervals for the trend estimate; the values represent the 2.5% and 97.5% percentiles of the posterior distribution of trend estimates (Sauer et al. 2017). Trends for which credible intervals do not include zero can be considered significant (red text represents a significant negative trend). "n/a" is used where there are data deficiencies or where the species does not breed, meaning no data available or not applicable.

| Region | Official Species Status | State/ Provincial Conservation Status | BBS Sample size (n) | BBS Trend 1967-2015 (%/yr, 95% CI) | BBS Trend 2005-2015 (%/yr, 95% CI) | Population estimate | Percentage of breeding population |
|---------------------------|-------------------------|---------------------------------------|---------------------|---------------------------------------|---------------------------------------|----------------------|-----------------------------------|
| Range wide | | | | | | | |
| Survey wide | | | 117 | -5.9 (-9.1, -2.8)^b | -2.8 (-7.6, 2.8) | 600,000 ^d | |
| United States | | | 61 | -0.7 (-3.3, 1.8) | 0.9 (-4.3, 6.6) | 400,000 | 75.5 ^d |
| Canada | T | | 56 | -8.0 (-11.6, -4.1)^b | -7.8 (-13.8, 2.01) | 130,000 | 32.5 |
| BCR-level | | | | | | | |
| Prairie Potholes BCR | | | 69 | -7.1 (-10.4, -3.5)^b | -5.0 (-10.8, 3.1) | 300,000 ^d | 45.8 |
| Shortgrass Prairie BCR | | | 8 | 1.4 (-3.3, 6.3) | 0.9 (-5.2, 6.7) | 110,000 | 19.5 |
| Badlands and Prairies BCR | | | 20 | -1.0 (-5.4, 3.3) | -0.4 (-10.3, 9.2) | 80,000 | 14.1 |
| Northern Rockies BCR | | | 20 | -2.7 (-6.8, 1.7) | -0.8 (-6.7, 9.1) | 110,000 | 20.2 |
| United States | | | | | | | |
| Arizona | | Tier 1C | n/a | n/a | n/a | n/a | n/a |
| Colorado | | Tier 2 | 8 | 1.4 (-3.3, 6.3) | 0.9 (-5.2, 6.7) | 80,000 | 13.7 |
| Kansas | | Tier 2 | n/a | n/a | n/a | n/a | n/a |
| Montana | | Tier 3 | 24 | -1.4 (-4.6, 1.8) | 0.2 (-8.1, 9.6) | 190,000 | 34.2 |
| Nebraska | | Tier 1 | n/a | n/a | n/a | n/a | n/a |
| New Mexico | | SGCN | n/a | n/a | n/a | n/a | n/a |
| North Dakota | | Tier 3 | n/a | n/a | n/a | n/a ^e | n/a |
| Oklahoma | | Tier 2 | n/a | n/a | n/a | n/a | n/a |
| Texas | | Tier S4 | n/a | n/a | n/a | n/a | n/a |
| Wyoming | | Tier 2 | 29 | -0.2 (-4.6, 4.4) | 1.1 (-4.5, 8.8) | 160,000 | 27.8 |
| Canada | | | | | | | |
| Alberta | | | 31 | -7.9 (-10.9, -5.1)^a | -9.6 (-18.2, -3.8)^a | 70,000 | 11.9 |
| Saskatchewan | | | 25 | -8.0 (-12.1, -3.0)^b | -7.4 (-14.4, 3.8) | 60,000 | 10.7 |

BBS trends: ^a High confidence, ^b Medium confidence, ^c Low confidence in reliability of the trend assessments (Sauer et al. 2017)

^d Population estimate and percentage of population are based on BBS data and other estimators, thus numbers and percentages by country, BCR, or state/province do not necessarily add up to the global population estimate or 100%, respectively (Blancher et al. 2013).

^e State estimates (95% CI) for McCown's Longspur in North Dakota in 1967, 1992, and 1993 were 50,000 (0-99,000), 4,000 (0-12,000), and 2,000 (0-6,000) breeding pairs respectively (Igl et al. 1999).